

Interactions between a surface-active cationic 3H-indole molecular probe and β -cyclodextrin. Design of a novel type of rotaxane

Xinghai Shen¹, Michel Belletête, Gilles Durocher^{*}

Département de Chimie, Université de Montréal, C.P. 6128, Succ. A, Montréal, Québec, Canada H3C 3J7

Received 14 September 1998; in final form 20 November 1998

Abstract

We report herein the interactions of a cationic surface-active molecular probe having long aliphatic chains, i.e., iodo-methyldioctadecyl 2-(*p*-hexylaminophenyl)-3,3-dimethyl-5-carboethoxy-3H-indole ammonium, with β -CD investigated by spectral and photophysical characterizations. It is found through lifetime measurements that only two species exist within the whole range of β -CD concentrations. Both the steady-state and the time-resolved fluorescence results further show that the stoichiometry of the inclusion complex is 1:3. It is also suggested that an interaction of the aliphatic chains of the cationic 3H-indole with β -CD takes place. Finally it is shown that a new rotaxane forms spontaneously in solution. © 1999 Elsevier Science B.V. All rights reserved.

1. Introduction

Cyclodextrins (CDs) are toroidally shaped cyclic oligosaccharides, mostly consisting of six, seven and eight glucose units for α -CD, β -CD and γ -CD, respectively. Their hydrophobic cavities enable them to accommodate various kinds of molecules to form inclusion complexes, which leads to widespread applications in pharmaceutical chemistry, food technology, analytical chemistry, chemical synthesis, and catalysis [1–6]. Therefore the investigations of the inclusion complexes have been the focus of great efforts in organic chemistry. Among the various

possibilities of intermolecular interactions between CDs and organic molecules, the 1:1 and 1:2 (guest:host) inclusion complexes are the most common types. However, under appropriate conditions, supramolecular assemblies such as catenanes [7], rotaxanes [8,9], polyrotaxanes [10], nanotubular structures [11,12], or threaded cyclodextrins [13] that do not involve any covalent bonding between the cyclodextrin and the other molecule, can be obtained.

In the past few years, our research group has been focused on the study of some substituted 3H-indoles in microenvironments [14–17]. Very recently, our research group started a program of studying the complexation between substituted 3H-indoles and CDs [18–22]. It was found that 1:1 and 1:2 complexes are usually formed between 3H-indoles and

^{*} Corresponding author. E-mail: durocher@umontreal.ca

¹ Present address: Department of Technical Physics, Peking University, Beijing 100871, P.R. China.




Fig. 2. Absorption (A) and fluorescence (B) spectra (normalized according to the respective fluorescence quantum yield) of **2** in: (1) water (solid); (2) 0.004 M β -CD (dash); (3) 0.008 M β -CD (dot); and (4) 0.015 M β -CD (short dash).

tions are shown in Fig. 2. The optical characteristics of these spectra are compiled in Table 1.

Fig. 2 and Table 1 show that the absorption wavenumber does not change going from water to β -CD solutions. This might suggest that both the amino and indolic nitrogen are probably at the junctions of adjacent cyclodextrins and thus experience close contacts with water molecules. However, the FWHM values of the absorption and fluorescence

bands increase going from water to β -CD solutions, which is consistent with the blue shift observed in the fluorescence spectra. This indicates that **2** transfers from a polar to a less polar environment.

It is also noted from Table 1 that the fluorescence quantum yield of **2** increases with increasing the β -CD concentration. This strongly suggests that **2** moves from water to less aqueous sites that avoid the intramolecular twisting responsible for the stabilization of the TICT state and the quenching of the normal fluorescence [31,33,34]. But by comparing the fluorescence quantum yield value of **2** in 0.015 M β -CD solution with those of 2-(*p*-aminophenyl)-3,3-dimethyl-5-carboethoxy-3H-indole and 2-(*p*-dimethylaminophenyl)-3,3-dimethyl-5-carboethoxy-3H-indole in 0.004 M β -CD solution, which are 0.37 and 0.39, respectively [18], one can infer that in 0.015 M β -CD solution there are possibly some amount of **2** remaining in water.

3.2. Association constants

Since only one new species is formed in the presence of β -CD according to the lifetime measurements (see Section 3.3), the data have been analyzed according to our most recent paper [20].

The plot of I/I_0 vs. $[\text{CD}]_0$ is shown in Fig. 3A. The NLR analysis indicates that reasonable results (values of the variables, standard errors, 95% confidence intervals, correlation coefficient, and absolute sum of squares) can be obtained only when a 1:3 complex is formed. The fit converged well with a correlation coefficient $r^2 = 0.992$ (see Fig. 3A). The value of the association constant (K') estimated is

Table 1
Spectral characteristics of **2** in various environments

Medium	$\bar{\nu}_A^a$ (cm^{-1})	ϵ^b ($\text{M}^{-1} \text{cm}^{-1}$)	$\bar{\nu}_F^c$ (cm^{-1})	Stokes shift (cm^{-1})	FWHM _A (cm^{-1})	FWHM _F (cm^{-1})	Φ_F
Water (pH = 9.5)	25400	24200	20200	5200	4800	3000	0.013
0.004 M β -CD (pH = 9.5)	25300	(26600) ^d	20400	4900	(4800) ^d	3200	0.021
0.008 M β -CD (pH = 9.5)	25300	(22500) ^d	20600	4700	(5000) ^d	3300	0.065
0.015 M β -CD (pH = 9.5)	25400	(22000) ^d	20700	4700	(5100) ^d	3400	0.14

^aAbsorption wavenumber taken at the center of mass of the absorption band.

^bMolar absorption coefficient at the peak intensity maximum.

^cFluorescence wavenumber taken at the center of mass of the fluorescence band.

^dThese values, especially those at $[\beta\text{-CD}] = 8$ and 15 mM, are with errors due to the scattering in the absorption spectra resulting from the large size of the 1:3 complex [20].




Fig. 3. (A) Plot of the relative fluorescence intensity vs. $[CD]_0$ for **2** complexed to β -CD. The full line is the nonlinear regression fit to the experimental data points following eq. (12) of Ref. [20]. (B) $1/(I - I_0)$ as a function of $[CD]_0^{-3}$.

$$(5.0 \pm 0.6) \times 10^{-3}$$

fr(thrg)6572suppcomple then

CTAB form 1:1 and 1:2 complexes with β -CD (Ref. [19] and refs. cited therein). Thus, a similar phenomenon could occur for **2**, the surface active molecular probe. If the interaction between the aliphatic chains of **2** and β -CD does exist, then $[S(CD)_3]$ becomes an apparent concentration that does include the contribution from higher-order complexes (1:4...1:7) and the equilibrium involving the formation of the 1:3 complex will be moved toward the right. This is opposite to the effect of urea, which makes the equilibrium move toward the left. In the latter case, we have observed that the *apparent association constant* is reduced by the hydrophobic interaction between urea and 3H-indoles [18–20]. Obviously, in the former case, the *apparent association constant* of the 1:3 complex will increase to some extent. To shed some light on this problem, one can compare the association constants of **2** with those of **1** in the absence and presence of urea, respectively. The values for **1** without and with urea are $(3.7 \pm 0.4) \times 10^5$ and $(5.6 \pm 0.8) \times 10^4 \text{ M}^{-3}$, respectively [20]. The fact that the association constant of **2** is larger than that of **1** in the absence of urea is an indication that the aliphatic chains might fit into the β -CD cavity. On the other hand, in the presence of 3 M urea, the association constant of **2** is about four times higher than that of **1**. It is worth mentioning that the urea concentration used is very high such that the hydrophobic interaction of urea and the aliphatic chains is considerable. Moreover it was reported that the hydrophobic interaction of urea on the aliphatic chain of SDS exists and thus the CMC of SDS is markedly increased [18,35]. On the basis of these facts, we believe that the complexation of the aliphatic chains of **2** with β -CD takes place. Otherwise, the association constant value between **2** and β -CD would be obviously lower than that between **1** and β -CD.

It is important to point out here that the three β -CD molecules complexed with the aromatic chain are not undergoing dynamic exchange with free β -CD because of the rotaxane-like structural feature [20]. However, the β -CD molecules complexed with the aliphatic chains should have exchange with free β -CD. Since one or two β -CD molecules might accommodate each aliphatic chain, many types of higher-order complexes between **2** and β -CD such as 1:4, 1:5, 1:6 and 1:7 complexes could possibly exist.

3.3. Lifetime measurements

The fluorescence decay curve of **2** was measured in β -CD solutions of various concentrations. A global analysis was carried out by linking the fluorescence decay curves together. The results were judged by the statistical fitting parameters χ^2 for the individual single curve analysis and for the global analysis (χ_g^2). The statistical criteria to judge the quality of the fit include both graphical and numerical tests [20,21].

We attempted a global double exponential analysis, linking the decay curves together. Two lifetimes were obtained with a satisfactory χ_g^2 value (Table 2). The triple exponential analysis was also attempted on the same set of conditions. However, it did not bring about any improvement of χ_g^2 . It has to be pointed out here that the fluorescence decay of **2** in pure water analyzed individually is not well reproduced by a single exponential. A short lifetime of 0.28 ns with a normalized preexponential factor of 0.97 is obtained together with a longer lifetime of 1.13 ns with a preexponential factor of 0.03. This is the reason why when the decay in pure water is included in the global analysis at all concentrations of β -CD, the individual χ^2 in pure water is poor (see Table 2). A similar phenomenon was observed for **1** [20]. On the other hand, performing the global analysis at all concentrations of β -CD excluding the decay in pure water, one obtains results for B_1, τ_1 and B_2, τ_2 similar as those reported in Table 2. Therefore, we believe that only two species exist in all the samples except that in pure water. The component with the smaller lifetime should correspond to the free molecules of **2** in water whereas the component with the larger lifetime involves the inclusion complex. As discussed in Section 3.1, **2** is less in contact with water in the inclusion complex. Thus the quenching of the fluorescence observed in aqueous solutions is avoided [31,33,34], giving rise to higher values of the fluorescence quantum yield and lifetime for the inclusion complex.

Using the values of B_1 and B_2 at various concentrations of β -CD in Table 2, one can also estimate the association constant [20,21]. The plot of B_2/B_1 as a function of $[CD]_0^3$ should exhibit a straight line through the zero point [20], the slope of which is equal to the K' value. Fig. 4 illustrates this kind of




Fig. 4. Plot of the ratio of the preexponential factors (B_2/B_1) as a function of $[CD]_0^3$.

straight line with a correlation coefficient $r = 0.992$. The K' value is estimated to be $(1.3 \pm 0.1) \times 10^6 M^{-3}$, which is similar to that of **1** obtained through the similar method [20]. It is noted that this value is 2.6 times higher than that obtained from the NLR analysis on the fluorescence intensity. This kind of discrepancy also appears in the literature [20,36]. The possible reason is that the inclusion complex in the excited state is not exactly the same as that in the ground state. Nevertheless, the fact that the plot of B_2/B_1 vs. $[CD]_0^3$ indeed exhibits a straight line through the zero point strongly supports the formation of the 1:3 complex. Again, we found that the plots of B_2/B_1 against $[CD]_0$, $[CD]_0^2$ and $[CD]_0^4$, respectively, do not exhibit straight lines (figures not shown). Finally, it should be noted that, due to the discrepancy of the K' value, no information on the interactions between the aliphatic chains of **2** and β -CD can be obtained through the comparison of K' values of **1** and **2**.

4. Concluding remarks

The NLR analysis on the steady-state fluorescence intensity of **2** in β -CD suggests that only the model based on the formation of the 1:3 inclusion

complex is operative. The data of the lifetimes obtained by the global analysis reveal that there are only two discrete environments, i.e., bulk water and an inclusion complex. The analysis of the preexponential factors is in good agreement with the model of the 1:3 inclusion complex. A comparison of the association constants of **2** and **1** in the absence and presence of urea, respectively, also suggests that the two long aliphatic chains are also incorporated into β -CD cavities. It is also shown that the inclusion complex between **2** and β -CD forms a novel kind of rotaxane. To our knowledge, the spontaneous formation of rotaxanes in solution is reported for the first time in the literature.

Acknowledgements

We gratefully acknowledge the financial assistance of the National Sciences and Engineering Research Council of Canada and the "Fonds FCAR" (Québec) in the form of grants. We also thank Mr. Adrian Popowycz for the synthesis and purification of the substituted 3H-indole studied here. XS thanks Peking University, PR China, and University of Montreal, Canada, for the financial assistance provided by the exchange program.

References

- [1] M.L. Bender, M. Komiyama, *Cyclodextrin Chemistry*, Springer, New York, 1978.
- [2] W. Saenger, *Angew. Chem., Int. Ed. Engl.* 19 (1980) 344.
- [3] J. Szejtli, *Cyclodextrins and Their Inclusion Complexes*, Akadémiai Kiadó, Budapest, 1982.
- [4] S. Li, W.C. Purdy, *Chem. Rev.* 92 (1992) 1457.
- [5] J. Szejtli, *Cyclodextrin Technology*, Kluwer, Dordrecht, 1988.
- [6] G. Wenz, *Angew. Chem., Int. Ed. Engl.* 33 (1994) 803.
- [7] A. Luttringhaus, F. Cramer, H. Prinzbach, *Angew. Chem.* 69 (1957) 137.
- [8] R.S. Wylie, D.H. Macartney, *J. Am. Chem. Soc.* 114 (1992) 3136.
- [9] R. Isnin, A.E. Kaifer, *J. Am. Chem. Soc.* 113 (1991) 8188.
- [10] A. Harada, J. Li, M. Kamachi, *Nature (London)* 356 (1992) 325.
- [11] G. Li, L.B. McGown, *Science* 264 (1994) 249.
- [12] G. Pistolis, A. Malliaris, *J. Phys. Chem.* 100 (1996) 15562.
- [13] A. Harada, J. Li, M. Kamachi, *Nature (London)* 364 (1993) 516.
- [14] S. Nigam, M. Belletête, R.S. Sarpal, G. Durocher, *J. Chem. Soc., Faraday Trans.* 91 (1995) 2133.
- [15] S. Nigam, R.S. Sarpal, M. Belletête, G. Durocher, *J. Colloid Interface Sci.* 177 (1996) 143.
- [16] R.S. Sarpal, M. Belletête, G. Durocher, *J. Phys. Chem.* 97 (1993) 5007.
- [17] R.S. Sarpal, G. Durocher, *J. Photochem. Photobiol. A: Chem.* 80 (1994) 307.
- [18] X. Shen, M. Belletête, G. Durocher, *J. Phys. Chem. B* 101 (1997) 8212.
- [19] X. Shen, M. Belletête, G. Durocher, *Langmuir* 13 (1997) 5830.
- [20] X. Shen, M. Belletête, G. Durocher, *J. Phys. Chem. B* 102 (1998) 1877.
- [21] S. Nigam, G. Durocher, *J. Phys. Chem.* 100 (1996) 7135.
- [22] S. Nigam, G. Durocher, *J. Photochem. Photobiol. A: Chem.* 103 (1997) 143.
- [23] V.C. Reinsborough, S.W. Wetmore, *Langmuir* 11 (1995) 2476.
- [24] H. Mwakibete, R. Cristantino, D.W. Bloor, E. Wyn-Jones, J.F. Holzwarth, *Langmuir* 11 (1995) 57.
- [25] H. Mwakibete, R. Cristantino, D.W. Bloor, E. Wyn-Jones, *Langmuir* 10 (1994) 3328.
- [26] J.W. Park, H.J. Song, *J. Phys. Chem.* 93 (1989) 6454.
- [27] J.W. Park, H.J. Song, *J. Inclusion Phenomena Mol. Recognit. Chem.* 17 (1994) 277.
- [28] P. Skrabal, J. Steiger, H. Zellinger, *Helv. Chim. Acta* 58 (1975) 800.
- [29] A. Popowycz, M.Sc. Thesis, University of Montreal, Montreal, PQ, 1991.
- [30] B. Zelent, T. Ganguly, L. Farmer, D. Gravel, G. Durocher, *J. Photochem. Photobiol. A: Chem.* 56 (1991) 165.
- [31] M. Belletête, R.S. Sarpal, G. Durocher, *Can. J. Chem.* 72 (1994) 2239.
- [32] J.M. Beechem, M. Ameloot, L. Brand, *Anal. Instrum.* 14 (1985) 379.
- [33] R.S. Sarpal, M. Belletête, G. Durocher, *Can. J. Chem.* 71 (1993) 1570.
- [34] M. Belletête, S. Nigam, G. Durocher, *J. Phys. Chem.* 99 (1995) 4015.
- [35] N.R. Choudhury, J.C. Ahluwalia, *J. Chem. Soc., Faraday Trans. 1* 77 (1981) 3119.
- [36] S.D. Feyter, J. van Stam, N. Boens, F.C. De Schryver, *Chem. Phys. Lett.* 249 (1996) 46.