



Original article

Synthesis of anionic sulphonate-crown ether surfactants



Jing Zhou, He-Le Liang, Qing-De Chen*, Xing-Hai Shen*

Beijing National Laboratory for Molecular Sciences (BNLMS), Radiochemistry and Radiation Chemistry Key Laboratory of Fundamental Science, College of Chemistry and Molecular Engineering, Peking University, Beijing 100871, China

ARTICLE INFO

Article history:

Received 22 April 2013

Received in revised form 3 June 2013

Accepted 7 June 2013

Available online 16 August 2013

Keywords:

Crown ether

Sulphonate

Surfactant

Synthesis

ABSTRACT

Two sulphonate-crown ether surfactants (sodium (4'-dodecanoyldibenzo-18-crown-6)-4''(5'')-sulphonate, ammonium (4'-dodecanoyldibenzo-18-crown-6)-4''(5'')-sulphonate) were designed and synthesized via a two-step continuous method. The products were purified and

conductivity and dynamic light scattering techniques.

fluorescence,

© 2013 Qing-De Chen and Xing-Hai Shen. Published by Elsevier B.V. on behalf of Chinese Chemical Society. All rights reserved.

1. Introduction

Crown ether surfactants have the ability of forming micelles and recognizing cations simultaneously, which distinguishes them from the normal surfactants, and thus have great potentials in several applications [1–5]. Up to now, to the best of our knowledge, almost all crown ether surfactants are non-ionic, in which a lipophilic long-chain alkyl group is attached onto a crown ether [6,7]. However, their low solubility in water and the obvious temperature effect (*i.e.*, cloud point) limited their applications in some fields. In this study, we designed and synthesized two kinds of surface-active sulphonate-crown ether to improve the solubility of the crown ether surfactants


7.8 mmol) was added slowly, and the reaction mixture was mechanically stirred for 3 h. Then, the hot solution was cooled and decomposed with an ice-water mixture. After stirring for 4 h, the resulting pale yellow precipitates were collected by filtration, washed with water until a neutral pH value was achieved, and dried in vacuum at room temperature. The reaction was monitored by thin-layer chromatography (silica thin-layer plate, petroleum ether/dichloromethane/ethanol = 1/1/0.2, v/v/v).

Sulfonation of 4'-dodecanoyl-DB18C6: The dried mixture of mono- and didodecanoyl-DB18C6 (3.28 g) was dissolved in anhydrous dichloromethane (20 mL). At -10°C , chlorosulfonic acid (2 mL, 30.4 mmol) was added dropwise. Later, the reaction mixture was warmed to room temperature gradually, and was stirred for 24 h. Then, the reaction mixture was cautiously poured into an ice-water mixture, the pH value was adjusted to 9 using a saturated sodium carbonate solution (or ammonia), and the mixture was stirred overnight. After being centrifuged, the aqueous phase was separated from the dichloromethane phase. The aqueous phase was lyophilized in vacuum to provide a solid powder, which was washed with hot petroleum ether to remove the residual reactant. The residue was extracted with ethanol at room temperature. The extracts were dried by vacuum rotary evaporation. After being washed by a small amount of cold water to remove the residual inorganic salts and dried in vacuum at room temperature, the final product was obtained. With respect to dichloromethane phase, didodecanoyl-DB18C6 was obtained after the removal of solvent and the following recrystallization from ethanol. The synthetic methods used to prepare the target anionic sulphonate-crown ether surfactants are outlined in Scheme 1.

g,

* Corresponding authors.

E-mail addresses: qdchen@pku.edu.cn (Q.-D. Chen),
xshen@pku.edu.cn (X.-H. Shen).



Scheme 1. Synthetic route of anionic sulphonate-crown ether surfactants: (a) lauric acid, PPA; (b) HSO₃Cl, CH₂Cl₂; (c) H₂O, Na₂CO₃/NH₃·H₂O.

2.2. Physicochemical characterization of the anionic sulphonate-crown ether surfactants

The fluorescence spectra were measured on an F-4500 Fluorescence Spectrometer (Hitachi, Japan). The excitation and emission slit widths were 10 nm and 5 nm, respectively, and the scanning speed was 240 nm/min. Conductivity measurements were carried out at 25.0 °C using a FE30 conductivity meter (Mettler Toledo, Switzerland). The conductivity meter was calibrated by measuring the conductivity of 1.0×10^{-2} mol L⁻¹ potassium chloride. The dynamic light scattering (DLS) experiment was carried out at 25.0 °C via a Zetasizer Nano ZS90 (Malvern Instruments, U.K.) laser light scattering spectroscopy (wavelength: 633 nm) with a scattering angle of 90°. The intensity autocorrelation functions were analyzed to obtain the hydrodynamic radii (R_h) of micelles with methods of Contin and Cumulant.

3. Results and discussion

3.1. Synthesis of anionic sulphonate-crown ether surfactants

In the acylation of benzo-crown ether, carboxylic acid-PPA is always used [8,9]. According to Grebenyuk's method [9], DB18C6 was acylated. However, the results of ¹H NMR analysis showed that the ratio of hydrogen atoms in the crown ether to those in the long chain was between 16:23 and 16:46, indicating that the products were a mixture of mono- and didodecanoyl-DB18C6. Different feed ratios of lauric acid to DB18C6 ranging from 1.1:1 to 1.9:1 were used. It was found that the increased feed ratio could not avoid the formation of didodecanoyl-DB18C6 while the lower feed ratio led to the residue of DB18C6 as well as the generation of didodecanoyl-DB18C6. The feed ratio of 1.3:1 was found to be proper to ensure the complete consumption of DB18C6. Due to the extremely similar properties of mono- and didodecanoyl-DB18C6, it was difficult to separate them. To solve this problem, we used a two-step continuous method, i.e., the neutralized and dried acylated product was directly used in the second step without any separation. After sulfonation, the segregation of didodecanoyl-DB18C6 from the target product became much easier.

In addition, it is worthwhile to notice that the usage of chlorosulfonic acid is necessary. Sulfate salt-PPA is a mild sulfonating reagent in the sulphonation of benzo-crown ether [10]. Nevertheless, in our work, when Na₂SO₄-PPA was used, the FTIR spectra of the product showed that the absorption peaks at ca. 2920 cm⁻¹ and 2850 cm⁻¹, corresponding to the stretching vibration of C–H group, were extremely weak, suggesting that the dodecanoyl group was lost at the reaction temperature of 75 °C. While concentrated sulfuric acid was tried, the dodecanoyl group was still lost even the reaction temperature was maintained –20 °C. Thus, chlorosulfonic acid was used to sulphonylate the monododecanoyl-DB18C6, and

then the product was hydrolyzed in a basic solution to produce the desired sulphonate. Using this strategy, the sulphonated products were finally obtained.

The products were characterized by IR, ¹H NMR and elemental analysis and the following data agree well with the structure of sodium (4'-dodecanoyldibenzo-18-crown-6)-4''(5'')-sulphonate and ammonium (4'-dodecanoyldibenzo-18-crown-6)-4''(5'')-sulphonate.


Sodium (4'-dodecanoyldibenzo-18-crown-6)-4''(5'')-sulphonate: Yellow powder, 0.86 g (22.2%), IR (KBr, cm⁻¹): ν 2924 (CH₂, ν_a(CH₂)), 2852 (CH₂, ν_s(CH₂)), 1672 (ArCO, ν_{C=O}), 1595 (Ar, ν_{carbon skeleton}), 1514 (Ar, ν_{carbon skeleton}), 1456 (Ar, ν_{carbon skeleton}), 1431 (COCH₂, δ_{CH₂}), 1267 (SO₃Na, ν_a(SO₂)), 1138 (CH₂OCH₂, ν_a(C–O–C)), 1041 (SO₃Na, ν_s(SO₂)), 694 (SO₃Na, ν_i(S–O)). ¹H NMR (400 MHz, DMSO-*d*₆): δ 0.849 (t, 3H, CH₃), 1.235 (m, 16H, CH₂), 1.578 (q, 2H, COCH₂CH₂), 2.953 (t, 2H, COCH₂), 3.917 (m, 8H, ArOCH₂CH₂), 4.109 (m, 4H, ArOCH₂), 4.196 (m, 4H, ArOCH₂), 6.913 (d, 1H, ArH), 7.085 (d, 1H, ArH), 7.163 (s, 1H, ArH), 7.185 (s, 1H, ArH), 7.451 (d, 1H, ArH), 7.647 (d, 1H, ArH). Elem. anal. calcd. % for C₃₂H₄₅NaO₁₀S: C 59.61, H 7.03; found: C 59.55, H 7.17.

Ammonium (4'-dodecanoyldibenzo-18-crown-6)-4''(5'')-sulphonate: Yellow powder, 0.91 g (23.5%), IR (KBr, cm⁻¹): ν 2924 (CH₂, ν_a(CH₂)), 2852 (CH₂, ν_s(CH₂)), 1674 (ArCO, ν_{C=O}), 1593 (Ar, ν_{carbon skeleton}), 1512 (Ar, ν_{carbon skeleton}), 1456 (Ar, ν_{carbon skeleton}), 1429 (COCH₂, δ_{CH₂}), 1267 (SO₃NH₄, ν_a(SO₂)), 1138 (CH₂OCH₂, ν_a(C–O–C)), 1063 (SO₃NH₄, ν_s(SO₂)). ¹H NMR (400 MHz, DMSO-*d*₆): δ 0.850 (t, 3H, CH₃), 1.237 (m, 16H, CH₂), 1.578 (q, 2H, COCH₂CH₂), 2.939 (t, 2H, COCH₂), 3.857 (m, 8H, ArOCH₂CH₂), 4.146 (m, 8H, ArOCH₂), 7.040 (d, 1H, ArH), 7.083 (d, 1H, ArH), 7.181 (s, 1H, ArH), 7.367 (s, 1H, ArH), 7.428 (d, 1H, ArH), 7.610 (d, 1H, ArH). Elem. anal. calcd. (%) for C₃₂H₄₇NO₁₀S·1H₂O: C 58.61, H 7.53, N 2.14; found: C 58.83, H 7.52, N 2.19.

3.2. Physicochemical properties of the anionic sulphonate-crown ether surfactants

It was found that the two kinds of synthesized surfactants had good solubility in ethanol and dimethyl sulfoxide (DMSO), but were difficult to be dissolved in dichloromethane. The solubility of the ammonium salt of the surfactants in water was lower than 5.0×10^{-5} mol L⁻¹, while their sodium salt can reach 2.0×10^{-3} mol L⁻¹ at least. This phenomenon indicates that the cations have a great impact on the solubility of the synthesized anionic sulphonate-crown ether surfactants. Therefore, the physicochemical characterization was focused on the sodium salt. Moreover, with the increase of the temperature, a cloud point phenomenon in the aqueous solution of the sodium salt was not observed.

When the ethanol solution of the sodium salt was excited at 322 nm, an obvious fluorescence emission peak appeared at ca.



395 nm (curve m in Fig. 1). When water was used as the solvent, the fluorescence emission peak was red-shifted to ca. 404 nm (curves c–l in Fig. 1). With the increase of the concentration of the sodium salts in aqueous solution, the fluorescence emission was enhanced, which could be ascribed to the aggregation of the surfactant and the formation of micelles. The plots of the relative fluorescence intensity vs the concentration of sodium salt (inset, Fig. 1) shows that a significant inflexion appears at the concentration of $8.5 \times 10^{-5} \text{ mol L}^{-1}$, which divides the plots into two linear regions. It can be concluded that the critical micelle concentration (*cmc*) of the surfactant is about $8.5 \times 10^{-5} \text{ mol L}^{-1}$. The results of the conductivity measurements (Fig. 2) indicated that the *cmc* of the surfactant is about $7.8 \times 10^{-5} \text{ mol L}^{-1}$, close to the result of fluorescence analysis.

Fig. 3 shows the DLS results of size distributions of the aqueous solution of the sodium salt. When the concentration of the sodium salt is 0.08 mmol L^{-1} , close to the *cmc* value, there already exist two peaks. The first peak ($R_h = 25 \text{ nm}$) is assigned to the formed small micelles, and the second peak ($R_h = 160 \text{ nm}$) is assigned to the formed large micelles. As the concentration of the sodium salt

increases to 0.10 mmol L^{-1} , the R_h value of the two kinds of micelles increase to 35 nm and 190 nm, respectively.

4. Conclusion

A novel series of anionic sulphonate-crown ether surfactants, *i.e.*, sodium (4'-dodecanoyldibenzo-18-crown-6)-4''(5'')-sulphonate and ammonium (4'-dodecanoyl-dibenzo-18-crown-6)-4''(5'')-sulphonate, were successfully synthesized in high purity in this work. The structures of these surfactants were elucidated by IR, ^1H NMR and elemental analysis. The experimental results indicated that the sodium salt was temperature insensitive, and had improved water solubility than nonionic crown ether surfactants. These properties allow them to have potential applications in several fields.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Nos. 91226112 and 20871009). We thank Professor Wen-Ting Hua from Peking University, for the generous help in the synthesis and purification of products.

References

- [1] G.W. Gokel, W.M. Leevy, M.E. Weber, Crown ethers: sensors for ions and molecular scaffolds for materials and biological models, *Chem. Rev.* 104 (2004) 2723–2750.
- [2] F.F. Chou, J.S. Shih, Micellar formation study of crown ether surfactants, *J. Chin. Chem. Soc.* 49 (2002) 599–605.
- [3] A. Gobbi, D. Landini, A. Maia, et al., Macrocyclic polyethers as enolate activators in base-catalyzed phase-transfer reactions, *J. Org. Chem.* 63 (1998) 5356–5361.
- [4] W.M. Leevy, M.E. Weber, M.R. Gokel, et al., Correlation of bilayer membrane cation transport and biological activity in alkyl-substituted lariat ethers, *Org. Biomol. Chem.* 3 (2005) 1647–1652.
- [5] L.L. Cui, D.M. Chao, J.F. Zhang, et al., Synthesis and properties of novel electro-active polyamide containing crown ether in the main chain, *Synth. Met.* 160 (2010) 400–404.
- [6] N.J. Turro, P.L. Kuo, Novel excimer formation of pyrene in micelles of a crown-ether surfactant, *J. Phys. Chem.* 91 (1987) 3321–3325.
- [7] K. Sabah, T. Heidelberg, R. Hashim, Novel crown ethers on glucose based glycolipids, *Carbohydr. Res.* 346 (2011) 891–896.
- [8] B. Gourdet, K. Singh, A.M. Stuart, et al., D1(1H, 1H, 2H, 2H-perfluorooctyl)-dibenzo-18-crown-6: a “light fluoruous” recyclable phase transfer catalyst, *J. Fluor. Chem.* 131 (2010) 1133–1143.
- [9] A.D. Grebenyuk, V.I. Vinogradova, A.K. Tashmukhamedova, Reaction of cytosine with 4'-substituted dibenzo-18-crown-6 sulfonylchlorides, *Chem. Nat. Compd.* 38 (2002) 182–185.
- [10] A.D. Grebenyuk, A.K. Tashmukhamedova, Sulfonation of 4'-acetyl-4'-(tert-butyl)-dibenzo-18-crown-6 and (dibenzo-18-crown-6)-4'-sulfonic acid with potassium sulfate in polyphosphoric acid, *Chem. Heterocycl. Compd.* 45 (2009) 297–301.